

- normal, obese and obese-diabetic subjects. *Metabolism* 2001; 50: 573-82.
15. Stumvoll M, Van Haefen T, Fritsche A, Gerich J. Oral glucose tolerance test indexes for insulin sensitivity and secretion based on various availabilities of sampling times. *Diabetes Care* 2001; 24: 796-7.
 16. Sainaghi PP, Castello L, Bergamasco L, et al. Metabolic characteristics of glucose intolerance: the critical role of obesity. *Exp Clin Endocrinol Diabetes*. 2008; 116: 86-93.
 17. Tam LS, Tomlinson B, Chu TT, et al. Cardiovascular risk profile of patients with psoriatic arthritis compared to controls--the role of inflammation. *Rheumatology (Oxford)*. 2008; 47: 718-23.
 18. Manrique-Arija S, Ureña I, Valdivielso P, et al. Insulin resistance and levels of adipokines in patients with untreated early rheumatoid arthritis. *Clin Rheumatol*. 2016; 35: 43-53.
 19. Labitigan M, Bahçe-Altuntas A, Kremer JM, et al. Higher rates and clustering of abnormal lipids, obesity, and diabetes mellitus in psoriatic arthritis compared with rheumatoid arthritis. *Arthritis Care Res (Hoboken)*. 2014; 66: 600-7.
 20. Castañeda S, Martín-Martínez MA, González-Juanatey C, et al. Cardiovascular morbidity and associated risk factors in Spanish patients with chronic inflammatory rheumatic diseases attending rheumatology clinics: Baseline data of the CARMA Project. *Semin Arthritis Rheum*. 2015; 44: 618-26.
 21. Burska AN, Sakthiswary R, Sattar N. Effects of tumour necrosis factor antagonists on insulin sensitivity/resistance in rheumatoid arthritis: a systematic review and meta-analysis. *PLoS One* 2015; 10: e0128889.

Non-commercial use only