

**Pain in systemic sclerosis**

Stefano Stisi,<sup>1</sup> Piercarlo Sarzi-Puttini,<sup>2</sup> Maurizio Benucci,<sup>3</sup> Giovanni Biasi,<sup>4</sup> Salvatore  
Bellissimo,<sup>1</sup> Rossella Talotta,<sup>2</sup> Fabiola Atzeni<sup>2</sup>

<sup>1</sup>Unit of Rheumatology, A.O. G. Rummo, Benevento; <sup>2</sup>Unit of Rheumatology, L. Sacco  
University Hospital, Milan; <sup>3</sup>Unit of Rheumatology, Ospedale San Giovanni di Dio, Florence;

<sup>4</sup>Unit of Rheumatology, University of Siena, Italy

**Correspondence:** Stefano Stisi, Unit of Rheumatology, A.O. G. Rummo, Benevento, Italy.

E-mail: [stefano.stisi@ao-rummo.it](mailto:stefano.stisi@ao-rummo.it)

**Key words:** *Spondyloarthritis, fibromyalgia, Prevalence, Disease activity, Ankylosing  
spondylitis disease activity score, Bath ankylosing spondylitis disease activity index.*

This paper must be considered as retracted due to a plagiarism misconduct.

See the *Retraction note* at: <https://doi.org/10.4081/reumatismo.2018.1171>